

Semigroup Forum (2011) 83:313–319
 DOI 10.1007/s00233-011-9311-0

RESEARCH ARTICLE

On regularity of sup-preserving maps: generalizing Zareckiĭ's theorem

Ulrich Höhle · Tomasz Kubiak

Received: 7 March 2011 / Accepted: 13 May 2011 / Published online: 8 June 2011
 © The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract A sup-preserving map f between complete lattices L and M is regular if there exists a sup-preserving map g from M to L such that $fgf = f$. In the class of completely distributive lattices, this paper demonstrates a necessary and sufficient condition for f to be regular. When $L = M$ is a power set, our theorem reduces to the well known Zareckiĭ's theorem which characterizes regular elements in the semigroup of all binary relations on a set. Another application of our result is a generalization of Zareckiĭ's theorem for quantale-valued relations.

Keywords Sup-preserving map · Regular map · Complete distributivity · Regular relation · Quantale · Quantale-valued relation

1 Introduction

A morphism $A \xrightarrow{f} B$ in a category \mathbf{C} is *regular* if there is a morphism $B \xrightarrow{g} A$ in \mathbf{C} with $fgf = f$ (cf. [4]).

The purpose of this paper is to examine regular morphisms in the category **Sup** whose objects are all complete lattices and whose morphisms are all sup-preserving

Communicated by Boris M. Schein.

The second named author gratefully acknowledges the grant MTM2009-12872-C02-02 from the Ministry of Science and Innovation of Spain.

U. Höhle
 Fachbereich C Mathematik und Naturwissenschaften, Bergische Universität, Gaußstraße 20,
 42097 Wuppertal, Germany
 e-mail: uhohhle@uni-wuppertal.de

T. Kubiak (✉)
 Wydział Matematyki i Informatyki, Uniwersytet im. Adama Mickiewicza, Umultowska 87,
 61-614 Poznań, Poland
 e-mail: tkubiak@amu.edu.pl

maps (cf. [3]). We give a sufficient condition for a morphism of **Sup** to be regular as well as a necessary one. Both these conditions involve complete distributivity, so that—when gathered together—they provide a characterization of regular morphisms of the category of all completely distributive lattices and their sup-preserving maps (see Theorem 3.1). Our result can be viewed as a non-atomic generalization of the famous Zareckiĭ's theorem which characterizes regular elements in the semigroup of all binary relations on a set (cf. [14]). As a matter of fact, we generalize a recent extension of Zareckiĭ's theorem stated in [12] which characterizes regular morphisms in the category of all relations. We recall that $\rho \subseteq X \times Y$ is called *regular* if there exists $\sigma \subseteq Y \times X$ such that $\rho\sigma\rho = \rho$. The result of [12] says: *A relation $\rho \subseteq X \times Y$ is regular iff the lattice $\{\rho(A) \mid A \subseteq X\}$ is completely distributive* (see Sect. 2 for notation if needed and cf. Note 1.2 below). In the case $X = Y$ we have the just mentioned Zareckiĭ's theorem. It may be remarked that with an idempotent $\rho \subseteq X \times X$ (i.e. $\rho\rho = \rho$), the Zareckiĭ's theorem is already seen in [9].

Another application of Theorem 3.1 is a generalization of Zareckiĭ's theorem for quantale-valued relations (see Theorem 4.3).

The extensive literature related to Zareckiĭ's theorem include [1, 5, 8, 11–13] among others. In particular, alternative proofs are given in [1, 11, 13] all of which use the fact that $(2^X, \subseteq)$ is an atomic completely distributive lattice (2^X is the power set of X).

Note 1.1 When saying that a morphism f of **Sup** is regular with $fgf = f$, we of course mean that g is a morphism of **Sup** too. We recall that in **Set** each map is regular (cf. [4]).

Note 1.2 Let $L \xrightarrow{f} M$ be a morphism in **Sup**. Then the range $f(L)$ is a complete lattice w.r.t. the partial ordering inherited from M and sups in $f(L)$ are formed in M , but in general not infs. Obviously, $f(L)$ together with the inclusion map is the image of f w.r.t. the (extremal epi, mono)-factorization property in **Sup**. In this context, saying that $f(L)$ is completely distributive simply means that the complete lattice $f(L)$ is completely distributive.

2 Binary relations as sup-preserving maps

Consider the category **Rel** whose objects are sets and $\rho : X \rightarrow Y$ is a morphism if $\rho \subseteq X \times Y$. The composition of $\rho \subseteq X \times Y$ and $\sigma \subseteq Y \times Z$ is the relation $\sigma\rho \subseteq X \times Z$ where

$$\sigma\rho = \{(x, z) \in X \times Z \mid \exists y \in Y : (x, y) \in \rho \text{ and } (y, z) \in \sigma\}.$$

Given $\rho \subseteq X \times Y$ and $A \subseteq X$, let $\rho(A)$ be defined by $\rho(A) = \bigcup_{x \in A} \rho(x)$ where $\rho(x) = \{y \in Y \mid (x, y) \in \rho\}$. The assignment $A \mapsto \rho(A)$ preserves arbitrary unions, hence $\{\rho(A) \mid A \subseteq X\}$ is a complete lattice w.r.t. the partial ordering inherited by 2^Y .

Thus, each $\rho \subseteq X \times Y$ determines a union-preserving map $2^X \xrightarrow{\Phi(\rho)} 2^Y$ by

$$\Phi(\rho)(A) = \rho(A).$$

Conversely (cf. [2, 6]), any map $2^X \xrightarrow{f} 2^Y$ determines $\Sigma(f) \subseteq X \times Y$ by

$$(x, y) \in \Sigma(f) \Leftrightarrow y \in f(\{x\}).$$

If f is union-preserving, then Φ sends $\Sigma(f)$ back to f (cf. property (2) below). The category **Rel** is isomorphic to a full subcategory of **Sup**. In fact, the following hold:

Proposition 2.1 *Let $\rho \subseteq X \times Y$ and $\sigma \subseteq Y \times Z$ be relations, let $2^X \xrightarrow{f} 2^Y \xrightarrow{g} 2^Z$ be arbitrary maps. Then:*

- (1) $\Phi(\sigma\rho) = \Phi(\sigma)\Phi(\rho)$ and $\Sigma(gf) = \Sigma(g)\Sigma(f)$,
- (2) $\Sigma(\Phi(\rho)) = \rho$, and $\Phi(\Sigma(f)) = f$ provided f is union-preserving.

From Proposition 2.1 we immediately obtain:

Fact 2.2 *The following statements hold:*

- (1) A relation $\rho \subseteq X \times Y$ is regular iff $2^X \xrightarrow{\Phi(\rho)} 2^Y$ is regular.
- (2) A union-preserving map $2^X \xrightarrow{f} 2^Y$ is regular iff $\Sigma(f) \subseteq X \times Y$ is a regular relation.

The Zareckiĭ's theorem (in its more general version of [12]) can now be formulated as follows:

Theorem 2.3 *A union-preserving map $2^X \xrightarrow{f} 2^Y$ is regular iff $f(2^X)$ is a completely distributive lattice.*

This formulation is an invitation to consider it in a more general lattice-theoretic setting by replacing the power sets by completely distributive lattices (see Theorem 3.1). We recall that the proofs of Zareckiĭ's theorem given in [1, 11, 13] make an essential use of the fact that 2^X is an *atomic* completely distributive lattice. The proof of Theorem 3.1 is valid for arbitrary completely distributive lattices and is thus different from all of them.

3 Regular and sup-preserving maps

It is well known that a complete lattice L is completely distributive iff L^{op} is. In particular, for each family $\{A_i\}_{i \in I}$ of subsets of L the following holds:

$$\bigvee_{i \in I} \bigwedge A_i = \bigwedge_{\varphi \in \prod_{i \in I} A_i} \left(\bigvee_{i \in I} \varphi(i) \right).$$

Given a complete lattice L , we write $a \triangleleft b$ iff, whenever $C \subseteq L$ and $b \leq \bigvee C$, there is $c \in C$ with $a \leq c$ (cf. [3, 9]). Then L is completely distributive iff \triangleleft has the approximation property:

$$a = \bigvee \{b \in L \mid b \triangleleft a\}$$

for each $a \in L$. For all $a, b, c, d \in L$ the following hold (cf. [3]):

- (\triangleleft_1) $a \triangleleft b$ implies $a \leq b$,
- (\triangleleft_2) $c \leq a \triangleleft b \leq d$ implies $c \triangleleft d$,
- (\triangleleft_3) $a \triangleleft b$ implies $a \triangleleft c \triangleleft b$ for some $c \in L$.

The following is our generalization of the Zareckiĭ's theorem.

Theorem 3.1 *Let L and M be completely distributive lattices, and let $L \xrightarrow{f} M$ be sup-preserving. Then f is regular iff $f(L)$ is a completely distributive lattice.*

Since both the *if* part and the *only if* part of Theorem 3.1 hold true in a more general setting, we wish to split the theorem into two propositions from which Theorem 3.1 follows immediately. In fact, the *only if* part is also valid for L a continuous lattice (cf. Remark 3.4).

Proposition 3.2 *Let L and M be complete lattices, and let $L \xrightarrow{f} M$ be sup-preserving. If both M and $f(L)$ are completely distributive, then f is regular.*

Proof For every $y \in M$ we define $A_y = \{c \in L \mid y \leq f(c)\}$. Then we put

$$G_y = \left\{ b \in L \mid f(b) = \bigwedge f(A_y) \right\}$$

where \bigwedge is performed in $f(L)$. Define a map $M \xrightarrow{g} L$ by

$$g(x) = \bigvee_{y \triangleleft x} \bigvee G_y$$

for all $x \in M$. Let $A \subseteq M$. By using (\triangleleft_1)–(\triangleleft_3), we have $y \triangleleft \bigvee A$ iff $y \triangleleft x$ for some $x \in A$ so that $g(\bigvee A) = \bigvee (\bigcup_{x \in A} \bigcup_{y \triangleleft x} G_y) = \bigvee_{x \in A} g(x)$, hence g is sup-preserving.

In order to verify $fgf = f$ we proceed as follows. As f is sup-preserving we have $\bigwedge f(A_y) = f(\bigvee G_y)$ and subsequently

$$\bigvee_{y \triangleleft f(a)} \bigwedge f(A_y) = f\left(\bigvee_{y \triangleleft f(a)} \bigvee G_y\right) = fgf(a), \quad a \in L.$$

Now we make use of the complete distributivity of $f(L)$ and obtain:

$$\bigwedge_{\substack{\varphi \in \prod_{y \triangleleft f(a)} f(A_y)}} \left(\bigvee_{y \triangleleft f(a)} \varphi(y) \right) = fgf(a).$$

Because of $\varphi(y) \in f(A_y)$ we have $y \leq \varphi(y)$, and the relation

$$f(a) = \bigvee_{y \triangleleft f(a)} y \leq fgf(a)$$

follows. On the other hand $y \triangleleft f(a)$ implies $y \leq f(a)$, and consequently $f(a) \in f(A_y)$. Because of $\bigwedge f(A_y) \leq f(a)$ the relation $f g f(a) \leq f(a)$ holds. \square

Proposition 3.3 *Let L and M be complete lattices, let $L \xrightarrow{f} M$ be sup-preserving. If f is regular and L is completely distributive, then $f(L)$ is completely distributive too.*

Proof Let $f g f = f$ where $M \xrightarrow{g} L$ is sup-preserving, and let $a \in L$. Then $g f(a) \in L$ and by complete distributivity of L we have

$$g f(a) = \bigvee \{b \in L \mid b \triangleleft g f(a)\}.$$

We now check that $b \triangleleft g f(a)$ in L implies $f(b) \triangleleft f(a)$ in $f(L)$. Indeed, if $f(a) \leq \bigvee f(C)$ for some $C \subseteq L$, then $g f(a) \leq \bigvee g f(C)$ and there is a $c \in C$ such that $b \leq g f(c)$. This yields $f(b) \leq f(c)$, so that $f(b) \triangleleft f(a)$. Consequently,

$$\begin{aligned} f(a) &= f g f(a) \\ &\leq \bigvee \{f(b) \mid f(b) \triangleleft f(a)\} \\ &\leq \bigvee \{y \in f(L) \mid y \triangleleft f(a)\} \\ &\leq f(a). \end{aligned}$$

We have proved that \triangleleft has the approximation property in $f(L)$. \square

Remark 3.4 We recall that a complete lattice L is *continuous* if $a = \bigvee \{b \in L \mid b \ll a\}$ for all $a \in L$, where $b \ll a$ iff, whenever $a \leq \bigvee A$ for some $A \subseteq L$, there exists a finite subset $D \subseteq A$ such that $b \leq \bigvee D$. Every completely distributive lattice is continuous (see [3]). It is not difficult to observe that Proposition 3.3 maintains its validity for the relation \ll and continuous lattices.

4 A quantalic version of Zareckiĭ's theorem

In what follows we extend our previous considerations in Sect. 2 from the two-valued to the many-valued setting. More specifically, we will replace maps $\mathbf{2}^X \rightarrow \mathbf{2}^Y$ by maps $L^X \rightarrow L^Y$ where the complete lattice L replaces the two point lattice $\mathbf{2}$. Here and elsewhere L^X is the set of all maps from X into L ordered pointwisely.

For this purpose we first recall that a triple $Q = (Q, \leq, \&)$ is called a *quantale* if (Q, \leq) is a complete lattice, $(Q, \&)$ is a semigroup, and $\&$ distributes over arbitrary sups in both variables (cf. [10]). We begin with the following example (cf. [6, 7]).

Example 4.1 (Quantale $\mathcal{Q}(L)$) Let L be a complete lattice. On the set $\mathcal{Q}(L)$ of all sup-preserving self-maps on L we define a partial ordering \leq in a pointwise way. Then $(\mathcal{Q}(L), \leq)$ is again a complete lattice. In particular, the sup of $\mathcal{S} \subseteq \mathcal{Q}(L)$ is given by $(\bigvee \mathcal{S})(a) = \bigvee_{\sigma \in \mathcal{S}} \sigma(a)$ for all $a \in L$. The composition

$$(\sigma_1 \& \sigma_2)(a) = \sigma_2(\sigma_1(a)), \quad a \in L,$$

induces a semigroup operation $\&$ on $\mathcal{Q}(L)$ so that the resulting triple $\mathcal{Q}(L) = (\mathcal{Q}(L), \leq, \&)$ becomes a quantale (in fact, a unital quantale—a detail not needed for our purposes).

Let \mathcal{Q} be an arbitrary quantale, and X and Y be sets. A \mathcal{Q} -valued relation $R : X \rightarrow Y$ is a map $X \times Y \xrightarrow{R} \mathcal{Q}$. The *composition* of \mathcal{Q} -valued relations $R : X \rightarrow Y$ and $S : Y \rightarrow Z$ is defined for all $x \in X$ and $z \in Z$ by

$$(SR)(x, z) = \bigvee_{y \in Y} R(x, y) \& S(y, z).$$

When $L = \mathbf{2}$, the category of all $\mathcal{Q}(L)$ -valued relations can be identified with **Rel** because $\mathcal{Q}(\mathbf{2})$ is just the two-point lattice.

Now we observe that each $\mathcal{Q}(L)$ -valued relation $R : X \rightarrow Y$ determines a sup-preserving map $L^X \xrightarrow{\Phi(R)} L^Y$ defined for all $f \in L^X$ and $y \in Y$ by

$$\Phi(R)(f)(y) = \bigvee_{x \in X} R(x, y)(f(x)).$$

On the other hand, each sup-preserving map $L^X \xrightarrow{\varphi} L^Y$ induces a $\mathcal{Q}(L)$ -valued relation $X \times Y \xrightarrow{\Sigma(\varphi)} \mathcal{Q}(L)$ defined for all $a \in L$, $x \in X$, and $y \in Y$ by

$$\Sigma(\varphi)(x, y)(a) = \varphi(a1_x)(y),$$

where

$$a1_x(z) = \begin{cases} a, & z = x, \\ \perp, & z \neq x \end{cases}$$

and \perp is the bottom element of L . The following provides a $\mathcal{Q}(L)$ -valued counterpart (in fact, a generalization from $\mathbf{2}$ to L) of Proposition 2.1.

Proposition 4.2 *Let $R : X \rightarrow Y$ and $S : Y \rightarrow Z$ be $\mathcal{Q}(L)$ -valued relations, and let $L^X \xrightarrow{\varphi} L^Y \xrightarrow{\psi} L^Z$ be arbitrary maps. Then:*

- (1) $\Phi(SR) = \Phi(S)\Phi(R)$ and $\Sigma(\psi\varphi) = \Sigma(\psi)\Sigma(\varphi)$,
- (2) $\Sigma(\Phi(R)) = R$, and $\Phi(\Sigma(\varphi)) = \varphi$ provided φ is sup-preserving.

Proof The proofs of (1) and the first part of (2) follows immediately from the definitions of all the involved compositions. Proving the second part of (2) requires a simple observation that each $f \in L^X$ has the following decomposition: $f = \bigvee_{x \in X} f(x)1_x$.

Indeed,

$$\begin{aligned}\Phi(\Sigma(\varphi))(f)(y) &= \bigvee_{x \in X} \varphi(f(x)1_x)(y) \\ &= \varphi\left(\bigvee_{x \in X} f(x)1_x\right)(y) \\ &= \varphi(f)(y)\end{aligned}$$

for all $f \in L^X$ and $y \in Y$. □

When L is completely distributive, then so is L^X . Using Theorem 3.1 and Proposition 4.2 (also cf. Note 1.2), we obtain our quantalic analogue of Zareckiĭ's theorem which when $L = \mathbf{2}$ yields the characterization of regularity of usual relations as given in Theorem 2.3.

Theorem 4.3 *Let L be a completely distributive lattice. Then a $\mathcal{Q}(L)$ -valued relation $R : X \rightarrow Y$ is regular iff $\Phi(R)(L^X)$ is a completely distributive lattice.*

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Bandelt, H.-J.: Regularity and complete distributivity. *Semigroup Forum* **19**, 123–126 (1980)
2. Erceg, M.A.: Functions, equivalence relations, quotient spaces and subsets in fuzzy set theory. *Fuzzy Sets Syst.* **3**, 75–92 (1980)
3. Gierz, G., Hofmann, K.H., Keimel, K., Lawson, J.D., Mislove, M., Scott, D.S.: *A Compendium of Continuous Lattices*. Springer, Berlin, Heidelberg, New York (1980)
4. Mac Lane, S.: *Categories for the Working Mathematician*. Springer, New York, Heidelberg, Berlin (1971)
5. Markowsky, G.: Idempotents and product representations with applications to semigroup of binary relations. *Semigroup Forum* **5**, 95–119 (1972)
6. Mulvey, C.J., Pelletier, J.W.: A quantisation of the calculus of relations. In: *CMS Proceedings*, vol. 13, pp. 345–360. Am. Math. Soc., Providence (1992)
7. Pelletier, J.W., Rosický, J.: Simple involutive quantales. *J. Algebra* **195**, 367–386 (1997)
8. Plemmons, R.J., West, M.T.: On the semigroup of binary relations. *Pac. J. Math.* **35**, 743–753 (1970)
9. Raney, G.N.: A subdirect-union representation of completely distributive lattices. *Proc. Am. Math. Soc.* **4**, 518–522 (1953)
10. Rosenthal, K.I.: *Quantales and Their Applications*. Pitman Research Notes in Mathematics, vol. 348. Longman Scientific & Technical, Longman House, Burnt Mill, Harlow (1996)
11. Schein, B.M.: Regular elements of the semigroup of all binary relations. *Semigroup Forum* **13**, 95–112 (1976)
12. Xu, X.-Q., Luo, M.-K.: Regular relations and normality of topologies. *Semigroup Forum* **72**, 477–480 (2006)
13. Yang, J.C.: A theorem on the semigroup of binary relations. *Proc. Am. Math. Soc.* **22**, 134–135 (1969)
14. Zareckiĭ, K.A.: The semigroup of binary relations. *Mat. Sb.* **61**, 291–305 (1963) (in Russian)